

Table 1 Percent mortality of *Cx. quinquefasciatus* larvae using crude extract of *R. madagaskariensis* leaves

Larval Instars	Concentration (%)	Percent Mortality (Mean \pm SE)		
		24h	48h	72h
First	0.1	30.67 \pm 1.25	45.33 \pm 0.82	53.33 \pm 0.94
	0.2	38.67 \pm 1.63	53.33 \pm 1.41	60.00 \pm 0.82
	0.3	49.33 \pm 2.05	62.67 \pm 0.00	66.67 \pm 0.94
	0.4	58.67 \pm 1.63	69.33 \pm 0.94	72.00 \pm 1.47
	0.5	65.33 \pm 0.80	73.33 \pm 0.65	78.67 \pm 0.00
Second	0.1	24.00 \pm 0.94	32.00 \pm 0.82	36.00 \pm 1.70
	0.2	34.67 \pm 2.05	40.00 \pm 1.63	42.67 \pm 0.65
	0.3	52.00 \pm 2.16	52.00 \pm 2.05	60.00 \pm 0.82
	0.4	56.00 \pm 1.70	61.33 \pm 0.82	66.67 \pm 1.25
	0.5	62.67 \pm 1.41	66.67 \pm 0.47	72.00 \pm 0.00
Third	0.1	22.67 \pm 0.00	29.33 \pm 0.80	36.00 \pm 1.47
	0.2	32.00 \pm 0.94	37.33 \pm 1.25	41.33 \pm 2.45
	0.3	49.33 \pm 0.00	53.33 \pm 1.41	57.33 \pm 0.00
	0.4	53.33 \pm 1.94	58.67 \pm 1.25	72.00 \pm 1.41
	0.5	56.00 \pm 1.63	62.67 \pm 0.00	74.67 \pm 0.00
Fourth	0.1	0.00 \pm 0.00	8.00 \pm 1.25	24.00 \pm 0.00
	0.2	2.67 \pm 1.25	9.33 \pm 0.47	30.67 \pm 0.00
	0.3	6.67 \pm 0.00	13.33 \pm 2.87	34.67 \pm 0.47
	0.4	9.33 \pm 0.47	17.33 \pm 1.45	36.00 \pm 0.82
	0.5	13.33 \pm 1.28	24.00 \pm 1.63	41.33 \pm 2.16