

Table 2 Probit analysis of larvicidal activity of *Murraya koenigii* hexane leaf extract isolated fractions against vector mosquitoes

Vector mosquitoes	<i>Aedes aegypti</i>						<i>Anopheles stephensi</i>						<i>Culex quinquefasciatus</i>					
	LC <sub>50</sub> (ppm)	95% CL LL UL		LC <sub>90</sub> (ppm)	95% CL LL UL		LC <sub>50</sub> (ppm)	95% CL LL UL		LC <sub>90</sub> (ppm)	95% CL LL UL		LC <sub>50</sub> (ppm)	95% CL LL UL		LC <sub>90</sub> (ppm)	95% CL LL UL	
A	95.65	88.86	104.65	210.23	186.97	243.90	101.51	94.16	111.39	213.81	190.24	247.84	71.61	66.48	77.33	188.20	168.20	216.99
B	151.68	134.55	178.78	281.38	240.06	348.06	185.79	154.31	248.44	378.39	298.82	539.13	122.16	111.06	138.44	246.88	215.03	295.45
C	196.69	165.60	254.55	350.37	283.66	475.85	298.73	215.47	579.35	586.19	402.02	1210.78	225.54	175.67	351.29	473.32	348.70	791.00
D	35.06	29.47	39.65	80.43	74.61	88.09	42.51	37.35	46.96	89.52	83.01	98.18	27.20	20.70	32.30	76.07	70.62	83.12
E	65.38	62.01	68.83	140.82	131.64	152.44	67.79	63.82	71.97	157.77	145.23	174.35	79.97	74.44	86.68	197.66	176.04	228.97
F	73.00	68.86	77.56	165.02	151.38	183.21	104.40	94.55	119.16	254.32	217.02	314.94	72.54	66.58	79.38	171.45	151.53	201.97

