

Table 1 Effect of *Ocimum gratissimum* powder on adult *Anopheles gambiae* mosquitoes

Weight (g)	Mortality at 6hours interval (%)					
	6	12	18	24	30	36
1.0	0.00±0.00 ^a	3.33±3.33 ^{ab}	6.67±3.33 ^a	36.67±6.67 ^b	93.33±6.67 ^d	100.00±0.00 ^b
2.0	0.00±0.00 ^a	10.00±0.00 ^b	16.67±8.82 ^b	36.67±6.67 ^b	100.00±0.00 ^d	100.00±0.00 ^b
3.0	0.00±0.00 ^a	13.33±3.33 ^{bc}	26.67±3.33 ^c	56.67±3.33 ^c	80.00±5.77 ^c	100.00±0.00 ^b
4.0	0.00±0.00 ^a	13.33±6.67 ^{bc}	26.67±3.33 ^c	43.33±3.33 ^{bc}	66.67±3.33 ^b	100.00±0.00 ^b
5.0	0.00±0.00 ^a	16.67±3.33 ^c	46.67±3.33 ^d	73.33±3.33 ^d	100.00±0.00 ^d	100.00±0.00 ^b
Control	0.00±0.00 ^a	0.00±0.00 ^a	0.00±0.00 ^a	0.00±0.00 ^a	0.00±0.00 ^a	0.00±0.00 ^a

Note: Each value is a mean ± standard deviation of four replicates; Values with different letters horizontally are significantly different at p≤0.05 Duncan's test