

Table 1 Immature developmental and survival rates of *Culex quinquefasciatus* mosquitoes exposed to different temperature regimens

Water Temperature (°C)	Duration of Immature Life Stage (Days)			Survivorship of Immature Life Stage (%)		
	Larval	Pupal	Aggregate	Larval	Pupal	Aggregate
28.00±0.02 (Control)	3.98±0.03 <sup>bc*</sup>	3.98±0.03 <sup>bc*</sup>	3.98±0.03 <sup>bc*</sup>	74.66±10.21 <sup>d<sub>a</sub>**</sup>	91.05±2.80 <sup>d<sub>b</sub></sup>	82.86±6.50 <sup>d</sup>
30.00	4.01±0.02 <sup>c</sup>	4.01±0.02 <sup>c</sup>	4.01±0.02 <sup>c</sup>	51.50±7.10 <sup>c<sub>a</sub></sup>	63.38±12.50 <sup>c<sub>b</sub></sup>	57.44±9.80 <sup>c</sup>
32.00	3.60±0.25 <sup>b</sup>	3.60±0.25 <sup>b</sup>	3.60±0.25 <sup>b</sup>	28.63±5.98 <sup>b<sub>a</sub></sup>	44.24±6.70 <sup>b<sub>b</sub></sup>	36.44±6.34 <sup>b</sup>
34.00	2.70±0.16 <sup>a</sup>	2.70±0.16 <sup>a</sup>	2.70±0.16 <sup>a</sup>	2.89±2.30 <sup>a<sub>a</sub></sup>	5.08±3.20 <sup>a<sub>a</sub></sup>	3.99±2.75 <sup>a</sup>

Note: \*Values followed by same superscript alphabets, in a column, are not significantly different at p = 0.05; \*\*Values followed by same subscript alphabets, in a row of larval and pupal survivorship, are not significantly different at p = 0.05