

Table 2 Seasonal distribution and IRDs of *An. gambiae* and *An. funestus* estimated from PSC in Panshekara Kumbotso LGA:

Months	<i>An. gamb s.l</i>	<i>An. gamb s.l</i> /room	<i>A.gamb s.l</i> /person	An. funestus	An. fun/ room	An. fun./ person
May-Jul						
Control	-	-	-	16	1.6	0.8
NET	-	-	-	17	1.7	0.8
NET + RAMBO	01	0.1	0.05	07	0.7	0.4
Aug-Oct						
C	151	15.1	7.6	133	13.3	6.6
N	10	1	0.5	17	1.7	0.9
NR	15	1.5	0.8	15	1.5	0.8
Nov-Jan						
C	-	-	-	06	0.6	0.3
N	-	-	-	03	0.3	0.2
NR	-	-	-	02	0.2	0.1
Feb-Apr						
C	-	-	-	02	0.2	0.1
N	-	-	-	02	0.2	0.1
NR	-	-	-	00	0.2	00

Note: C= Control, N= netting, NR= netting + Rambo. Relative abundance of *An. gambiae sl.* and *An. funestus* group collected in Panshekara between May 2010 – April 2011. *An. funestus* is preponderant and responsible for the biting in the dry season (May – July)