

Table 3 Susceptibility of immature forms of *Cx. quinquefasciatus* to bioactive culture filtrate isolates of *B. cereus*

Instar	Conc (mg/ml)	Percent mortality (\pm SE) at different exposure period		
		24 hrs	48 hrs	72 hrs
First	2.0	100.00 \pm 0.00	100.00 \pm 0.00	100.00 \pm 0.00
	1.5	87.00 \pm 1.15	89.33 \pm 0.88	93.00 \pm 0.57
	1.0	72.33 \pm 0.88	76.00 \pm 1.15	76.66 \pm 0.33
	0.5	54.66 \pm 0.66	58.00 \pm 0.57	60.33 \pm 0.88
	0.3	37.33 \pm 1.20	38.00 \pm 1.15	40.00 \pm 1.15
	Control	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00
Second	2.0	95.00 \pm 0.57	98.66 \pm 0.33	100.00 \pm 0.00
	1.5	80.00 \pm 0.57	83.66 \pm 0.66	84.66 \pm 0.66
	1.0	66.33 \pm 0.88	67.33 \pm 0.88	69.33 \pm 0.33
	0.5	46.00 \pm 0.57	48.33 \pm 0.33	49.00 \pm 1.15
	0.3	31.67 \pm 0.88	35.00 \pm 1.73	37.00 \pm 1.15
	Control	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00
Third	2.0	92.66 \pm 0.66	94.66 \pm 0.88	100 \pm 0.00
	1.5	80.00 \pm 0.57	81.33 \pm 0.33	83.66 \pm 0.33
	1.0	51.66 \pm 0.88	53.66 \pm 0.88	53.00 \pm 0.57
	0.5	37.33 \pm 0.33	38.00 \pm 0.57	41.33 \pm 0.88
	0.3	30.00 \pm 1.45	32.00 \pm 1.15	34.00 \pm 1.53
	Control	0.00 \pm 0.00	0.66 \pm 0.33	1.0 \pm 0.57
Fourth	2.0	91.33 \pm 0.88	93.66 \pm 0.88	100.00 \pm 0.00
	1.5	75.33 \pm 0.33	78.00 \pm 1.15	81.33 \pm 0.88
	1.0	49.33 \pm 1.33	51.33 \pm 0.88	51.00 \pm 0.00
	0.5	31.00 \pm 0.00	34.33 \pm 0.88	35.66 \pm 0.66
	0.3	22.00 \pm 1.15	23.00 \pm 1.15	28.00 \pm 1.15
	Control	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00

Note: M: Mortality, SE: Standard error